

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Commissioner
 US Department of Commerce
 United States Patent and Trademark
 Office, PCT
 2011 South Clark Place Room 524
 Arlington, VA 22202
 ETATS-UNIS D'AMERIQUE
 ETATS-UNIS D'AMERIQUE
 in its capacity as elected Office

Date of mailing (day/month/year) 27 October 2000 (27.10.00)	
International application No. PCT/NZ00/00031	Applicant's or agent's file reference 25635 MRB
International filing date (day/month/year) 17 March 2000 (17.03.00)	Priority date (day/month/year) 17 March 1999 (17.03.99)
Applicant PODILA, Gopi, Krishna et al	

1. The designated Office is hereby notified of its election made:



in the demand filed with the International Preliminary Examining Authority on:

10 October 2000 (10.10.00)



in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was

was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO
 34, chemin des Colombettes
 1211 Geneva 20, Switzerland

Facsimile No.: (41-22) 740.14.35

Authorized officer

Manu Berrod

Telephone No.: (41-22) 338.83.38

PCT INTERNATIONAL COOPERATION TREATY

9/936869

PCT

From the INTERNATIONAL BUREAU

NOTIFICATION OF THE RECORDING
OF A CHANGE(PCT Rule 92bis.1 and
Administrative Instructions, Section 422)

To:

WEST-WALKER, Gregory, James
A J Park
6th Floor
Huddart Parker Building
1 Post Office Square, P.O. Box 949
Wellington 6015
NOUVELLE-ZÉLANDEDate of mailing (day/month/year)
15 January 2002 (15.01.02)Applicant's or agent's file reference
25635 MRB

IMPORTANT NOTIFICATION

International application No.
PCT/NZ00/00031International filing date (day/month/year)
17 March 2000 (17.03.00)

1. The following indications appeared on record concerning:

☒ the applicant ☒ the inventor ☐ the agent ☐ the common representative

Name and Address

LIU, Jun-Jun
900 Champion Street #7
Houghton, MI 49931
United States of America

State of Nationality

CN

State of Residence

US

Telephone No.

Facsimile No.

Teleprinter No.

2. The International Bureau hereby notifies the applicant that the following change has been recorded concerning:

☐ the person ☐ the name ☒ the address ☐ the nationality ☒ the residence

Name and Address

LIU, Jun-Jun
Canadian Forest Service
Pacific Forest Centre
506 West Burnside Rd
Victoria, BC, V8T 1M
Canada

State of Nationality

CN

State of Residence

CA

Telephone No.

Facsimile No.

Teleprinter No.

3. Further observations, if necessary:

4. A copy of this notification has been sent to:

☒ the receiving Office ☐ the designated Offices concerned
☐ the International Searching Authority ☒ the elected Offices concerned
☐ the International Preliminary Examining Authority ☐ other:The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland

Authorized officer

Gabriele BAEHR

Facsimile No.: (41-22) 740.14.35

Telephone No.: (41-22) 338.83.38

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF THE RECORDING
OF A CHANGE(PCT Rule 92bis.1 and
Administrative Instructions, Section 422)

From the INTERNATIONAL BUREAU

To:

WEST-WALKER, Gregory, James
A J Park
6th Floor
Huddart Parker Building
1 Post Office Square, P.O. Box 949
Wellington 6015
NOUVELLE-ZÉLANDE

Date of mailing (day/month/year) 23 March 2001 (23.03.01)	IMPORTANT NOTIFICATION
Applicant's or agent's file reference 25635 MRB	
International application No. PCT/NZ00/00031	International filing date (day/month/year) 17 March 2000 (17.03.00)

1. The following indications appeared on record concerning:									
<input type="checkbox"/> the applicant	<input type="checkbox"/> the inventor <input checked="" type="checkbox"/> the agent <input type="checkbox"/> the common representative								
Name and Address BENNETT, Michael, Roy West-Walker Bennett Mobil on the Park 157 Lambton Quay Wellington New Zealand	<table border="1"> <tr> <td>State of Nationality</td> <td>State of Residence</td> </tr> <tr> <td colspan="2">Telephone No. 64-4-499-9058</td> </tr> <tr> <td colspan="2">Facsimile No. 64-4-499-9306</td> </tr> <tr> <td colspan="2">Teleprinter No.</td> </tr> </table>	State of Nationality	State of Residence	Telephone No. 64-4-499-9058		Facsimile No. 64-4-499-9306		Teleprinter No.	
State of Nationality	State of Residence								
Telephone No. 64-4-499-9058									
Facsimile No. 64-4-499-9306									
Teleprinter No.									
2. The International Bureau hereby notifies the applicant that the following change has been recorded concerning:									
<input type="checkbox"/> the person <input checked="" type="checkbox"/> the name <input checked="" type="checkbox"/> the address <input type="checkbox"/> the nationality <input type="checkbox"/> the residence									
Name and Address WEST-WALKER, Gregory, James A J Park 6th Floor Huddart Parker Building 1 Post Office Square, P.O. Box 949 Wellington 6015 New Zealand	<table border="1"> <tr> <td>State of Nationality</td> <td>State of Residence</td> </tr> <tr> <td colspan="2">Telephone No. 64-4-473 8278</td> </tr> <tr> <td colspan="2">Facsimile No. 64-4-472 3358</td> </tr> <tr> <td colspan="2">Teleprinter No.</td> </tr> </table>	State of Nationality	State of Residence	Telephone No. 64-4-473 8278		Facsimile No. 64-4-472 3358		Teleprinter No.	
State of Nationality	State of Residence								
Telephone No. 64-4-473 8278									
Facsimile No. 64-4-472 3358									
Teleprinter No.									
3. Further observations, if necessary:									
4. A copy of this notification has been sent to:									
<input checked="" type="checkbox"/> the receiving Office	<input type="checkbox"/> the designated Offices concerned								
<input type="checkbox"/> the International Searching Authority	<input checked="" type="checkbox"/> the elected Offices concerned								
<input checked="" type="checkbox"/> the International Preliminary Examining Authority	<input type="checkbox"/> other:								

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer J. Leitao Telephone No.: (41-22) 338.83.38
---	---

INTERNATIONAL SEARCH REPORT

In .tional Application No

ECT/GB 98/00939

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C12N15/82 C12N15/11 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94 10320 A (MOGEN INT ; SIJMONS PETER CHRISTIAAN (NL); GODDIJN OSCAR JOHANNES M) 11 May 1994 * see the whole document, esp. p.3 1.7-12 *	1,2,4-6, 10-13, 15,18,21
X	WO 97 04113 A (DANISCO ; POULSEN PETER (DK)) 6 February 1997 * see esp. p.1 - p.14 1.14 *	1,2,4-6, 8,10,11, 13,15, 18,21
A	WO 93 10251 A (MOGEN INT) 27 May 1993 see the whole document	1-21
A	WO 95 25787 A (UNIV RUTGERS ; MALIGA PAL (US)) 28 September 1995 see the whole document	1-21

-/--



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

30 June 1998

Date of mailing of the international search report

10/07/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Kania, T

INTERNATIONAL SEARCH REPORT

In International Application No

PT/GB 98/00939

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
T	<p>WO 97 30162 A (FORBIO RESEARCH PTY LTD ;TEASDALE ROBERT DIXON (AU); MOURADOV AIDY) 21 August 1997 see the whole document -----</p>	1-21

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/00939

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9410320	A	11-05-1994	AU 5420594 A	24-05-1994
			CA 2148451 A	11-05-1994
			EP 0666922 A	16-08-1995
WO 9704113	A	06-02-1997	AU 6614696 A	18-02-1997
			EP 0839203 A	06-05-1998
WO 9310251	A	27-05-1993	AU 2928492 A	15-06-1993
			CA 2121578 A	27-05-1993
			EP 0668921 A	30-08-1995
			HU 70264 A	28-09-1995
			JP 7500970 T	02-02-1995
WO 9525787	A	28-09-1995	US 5530191 A	25-06-1996
			AU 687544 B	26-02-1998
			AU 2198195 A	09-10-1995
			BR 9507180 A	14-10-1997
			CA 2186164 A	28-09-1995
			CN 1148869 A	30-04-1997
			EP 0805852 A	12-11-1997
			JP 9510615 T	28-10-1997
			PL 316435 A	06-01-1997
WO 9730162	A	21-08-1997	AU 1713297 A	02-09-1997

PLANTS HAVING MODIFIED REPRODUCTIVE CAPACITY

This invention relates to plants having modified reproductive capacity. In particular, it relates to a plant reproductive tissue specific promoter and its use in promoting transcription/expression of associated genes in plant reproductive tissue, including for the purpose of producing plants which have diminished reproductive capacity or which are sterile.

INTRODUCTION

It is desirable that the genetic basis of reproduction/flower development in plants be determined. Identification of genes involved in plant reproduction and/or flower development together with the regulatory elements which control their expression in reproductive tissue allows for modulation of the reproductive capacity of plants and specifically enables the production of reproductively null (sterile) plants. Identification of the regulatory elements involved further allows for the expression in reproductive tissue of genes which are heterologous to the plant where that is desirable.

The applicants have now identified and isolated such a reproductive tissue specific promoter which endogenously regulates expression of a peptide involved in the reproductive cycle of *Pinus radiata*. It is broadly towards this promoter, to its homologs in other plants and to its use in effecting expression of associated genes within the reproductive tissue of plants that the present invention is directed.

SUMMARY OF THE INVENTION

In a first aspect, the present invention provides a polynucleotide which has a nucleotide sequence of from nucleotides 1 to 1320 of Figure 2 and which has the ability, when operatively associated with a nucleotide sequence encoding a peptide, to promote transcription of that nucleotide sequence, or a polynucleotide which is a functionally equivalent variant thereof.

In a second aspect, the invention provides a plant reproductive tissue promoter which has a nucleotide sequence of from nucleotides 1 to 1320 of Figure 2, or a functionally equivalent variant thereof.

5 In a further aspect, the invention provides a DNA construct which comprises:

- (a) a polynucleotide having activity as a transcriptional promoter as described above;
- (b) an open reading frame polynucleotide coding for a peptide; and
- 10 (c) a termination sequence.

In yet a further aspect, the invention provides a DNA construct which comprises:

- (a) a promoter sequence as defined above;
- 15 (b) an open reading frame polynucleotide coding for a peptide; and
- (c) a termination sequence.

In each construct the open reading frame can be in a sense orientation, or an anti-sense orientation.

20

In one embodiment, the open reading frame polynucleotide encodes a peptide having the sequence of Figure 1.

In other embodiments, the open reading frame polynucleotide encodes a peptide which, when expressed in reproductive tissue of a plant, causes said plant's reproductive organs to abort, to redefine themselves as vegetative, or to stop development.

25

In still another embodiment, the open reading frame polynucleotide encodes a peptide which, when expressed in the reproductive tissue of a plant, causes cell death.

30

In yet another embodiment, the open reading frame polynucleotide encodes a peptide, which when expressed in reproductive tissue of a flowering plant, causes an alteration in the timing of flowering of said plant.

5 In a preferred form, the construct further includes:

(d) a selection marker sequence.

10 In a further aspect, the invention provides a transgenic plant cell which includes a construct as described above.

By "transgenic" as used herein, the applicants mean containing non-endogenous genetic material.

15 In another aspect, the invention provides a transgenic plant which includes a construct as described above.

20 In still another aspect, the invention provides a transgenic plant which contains a polynucleotide having activity as a transcriptional promoter as described above or a reproductive tissue promoter as described above, which plant has a reduced reproductive capacity.

It is particularly preferred that the plant be sterile.

25 Conveniently, in said plant said polynucleotide or promoter is operatively associated with a nucleotide sequence encoding a RNase.

The plant can be a coniferous plant, such as a coniferous plant of the *Pinus* genus, or a tree such as a member of the *Eucalyptus* genus.

30

It is particularly preferred that the transgenic plant be a member of a species selected from *Pinus radiata*, *Pinus taeda*, *Pinus elliotti*, *Pinus clausa*, *Pinus palustris*, *Pinus echinata*, *Pinus ponderosa*, *Pinus jeffrey*, *Pinus resinosa*, *Pinus rigida*, *Pinus banksiana*, *Pinus serotina*, *Pinus strobus*, *Pinus monticola*, *Pinus lambertiana*, *Pinus*

virginiana, *Pinus contorta*, *Pinus cariboea*, *Pinus pinaster*, *Pinus brutia*, *Pinus eldarica*, *Pinus coulteri*, *Pinus nigra*, *Pinus sylvestris*, *Pinus tecunumannii*, *Pinus keysia*, *Pinus oocarpa* and *Pinus maximowii*.

5 DESCRIPTION OF THE DRAWINGS

While the invention is broadly as defined above, it will be appreciated by those persons skilled in the art that it is not limited thereto and that it further includes embodiments of which the following description provides examples. In addition, the invention will be better understood through reference to the accompanying drawings in which:

Figure 1 shows the amino acid sequence of the reproductive peptide PrAG1, together with the nucleotide sequence coding therefor;

Figure 2 shows the sequence of the PrAG1 promoter, which is the focus of the present invention, isolated from *Pinus radiata*;

Figure 3 is an RNA gel blot analysis of PrAG1 mRNA accumulation in *Pinus radiata* organs. Twenty µg of total RNA from various organs was electrophoresed, blotted onto nylon membranes, and hybridized with 3'-terminal fragment of PrAG1 cDNA. Total RNA was isolated from immature male cone(M), immature female cone(F), vegetative shoot (V)s, needle (N) and stem (S). The 26S and 18S rRNA was used as control (bottom);

Figure 4 is a DNA gel blot analysis of *Pinus radiata* genomic DNA hybridized with the 3' terminal region of PrAG1. 20 µg genomic DNA was digested with BamHI (BA) and Bgl II (BG) EcoRI(E), HindIII (H), XhoI(X);

Figure 5 is a Reverse Transcription-Polymerase Chain Reaction (RT-PCR) analysis showing reproductive-organ specific expression of PrAG1. RT-PCR analysis was performed on total RNA isolated from different organs of radiata Pine: (M) immature male cone, (F) immature female cone, (Vs) vegetative shoot, (N) needle and (S) stem. PrAG1 genes were amplified with PrAG1 gene-specific oligonucleotides. Products

from the PCR reactions were electrophoresed, blotted, and hybridized with a labelled probe of PrAG1 specific fragment;

Figure 6 shows the construction of pRAGPR;

5

Figure 7. PCR analysis of transgenic tobacco lines. Genomic DNA (200 ng each) from controls and putative transgenic tobacco plants was used as template along with primers for NPTII gene. Lane PC is positive control (20 ng pRAGPR plasmid was used as a template); C1-C3 are control nontransformed tobacco plants; lanes 10 1a, 1b, 2, 3a, 3b, 3c, 4a, 4b are transgenic plants. The size of NPT II gene PCR product is indicated on the right;

Figure 8. Southern analysis of transgenic and control tobacco lines to confirm integration of PrAG1promoter-RNase gene cassette. Genomic DNA (20 ug each) was 15 digested with Hind III and Sac I enzymes, electrophoresed and transferred to a nylon membrane. ³²-P labeled RNase gene was used as a probe. The size of PrAG1promoter-RNase gene cassette is indicated on the right. Designation of control and transgenic plants is as indicated in Figure 7; and

20 Figure 9. Southern analysis of transgenic and control tobacco lines to confirm integration profiles of PrAG1promoter-RNase gene cassette. Genomic DNA (20 ug each) was digested with Hind III enzyme, electrophoresed and transferred to a nylon membrane. ³²-P labeled RNase gene was used as a probe. Sizes of DNA fragments hybridizing to the RNase gene probe are indicated on the right. Designation of 25 control and transgenic plants is as indicated in Figure 7.

DESCRIPTION OF THE INVENTION

30 As broadly outlined above, the applicants have identified a plant promoter which is involved in plant reproduction. The promoter, which was isolated from *Pinus radiata* is called herein the "PrAG1 promoter".

The nucleotide sequence of the PrAG1 promoter is given in Figure 2 from 35 nucleotides 1 to 1320. It will however be appreciated that the invention is not

restricted only to the polynucleotide having that specific nucleotide sequence. Instead, the invention also extends to functionally equivalent variants of that polynucleotide.

- 5 The phrase "functionally equivalent variants" recognises that it is possible to vary nucleotide sequence while retaining substantially equivalent functionality.

Variants can have a greater or lesser degree of homology as between the variant nucleotide sequence and the original.

10

- Polynucleotide sequences may be aligned, and percentage of identical nucleotides in a specified region may be determined against another sequence, using computer algorithms that are publicly available. Two exemplary algorithms for aligning and identifying the similarity of polynucleotide sequences are the BLASTN and FASTA algorithms. The BLASTN software are available on the NCBI anonymous FTP server (ftp://ncbi.nlm.nih.gov) under /blast/executables/. The BLASTN algorithm version 2.0.4 [Feb-24-1998], set to the default parameters described in the documentation of variants according to the present invention. The use of the BLAST family of algorithms, including BLASTN, is described at NCBI's website at URL <http://www.ncbi.nlm.nih.gov/BLAST/newblast.html>. The computer algorithm FASTA is available on the Internet at the ftp site <ftp://ftp.virginia.edu/pub/fasta/>. Version 2.0u4, February 1996, set to the default parameters described in the documentation and distributed with the algorithm, is also preferred for use in the determination of variants according to the present invention. The use of the FASTA algorithm is described in W. R. Pearson and D. J. Lipman, "Improved Tools for Biological Sequence Analysis", *Proc. Natl. Acad. Sci. USA* 85:2444-2448 (1988).
- 15
- 20
- 25

The following running parameters are preferred for determination of alignments and similarities using BLASTN that contribute to E values (as discussed below) and

- 30 percentage identity: Unix running command: blastall -p blastn -d emblddb -e 10 -G 1 -E 1 -r 2 -v 50 -b 50 -I queryseq -o results; and parameter default values:
- p Program Name [String]
 - d Database [String]
 - e Expectation value (E) [Real]

- G Cost to open a gap (zero invokes default behaviour) [Integer]
- E Cost to extend a gap (zero invokes default behaviour) [Integer]
- r Reward for a nucleotide match (blastn only) [Integer]
- v Number of one-line descriptions (V) [Integer]
- 5 -b Number of alignments to show (B) [Integer]
- i Query File [File In]
- o BLAST report Output File [File Out] Optional

10 The "hits" to one or more database sequences by a queried sequence produced by BLASTN, FASTA, or a similar algorithm, align and identify similar portions of sequences. The hits are arranged in order of the degree of similarity and the length of sequence overlap. Hits to a database sequence generally represent an overlap over only a fraction of the sequence length of the queried sequence.

15 The BLASTN and FASTA algorithms also produce "Expect" or E values for alignments. The E value indicates the number of hits one can "expect" to see over a certain number of contiguous sequences by chance when searching a database of a certain size. The Expect value is used as a significance threshold for determining whether the hit to a database, such as the preferred EMBL database, indicates true
 20 similarity. For example, an E value of 0.1 assigned to a hit is interpreted as meaning that in a database of the size of the EMBL database, one might expect to see 0.1 matches over the aligned portion of the sequence with a similar score simply by chance. By this criterion, the aligned and matched portions of the sequences then have a 90% probability of being the same. For sequences having an E value of
 25 0.01 or less over aligned and matched portions, the probability of finding a match by chance in the EMBL database is 1% or less using the BLASTN or FASTA algorithm.

30 It is also recognised that as the function of the polynucleotide is as a transcriptional promoter there are regions of the polynucleotide which are more critical to, and characteristic of, this function than others. An example are the TATA boxes at positions 280 to 286, 282 to 288 and 1015 to 1021 from the 5' end of the sequence. Therefore, polynucleotides which include these regions of the polynucleotide of Figure 2 and have equivalent transcriptional functionality are contemplated

variants, even where there is a lesser degree of homology elsewhere in the sequence.

5 According to one embodiment, "variant" polynucleotides, with reference to the polynucleotide of the present invention, preferably comprise sequences having the same number or fewer nucleic acids than the polynucleotide of the present invention and producing an E value of 0.01 or less when compared to the polynucleotide of the present invention. That is, a variant polynucleotide is any sequence that has at least a 99% probability of being the same as the
10 polynucleotide of the present invention, measured as having an E value of 0.01 or less using the BLASTN or FASTA algorithms set at the parameters discussed above.

Variant polynucleotide sequences will also generally hybridize to the recited polynucleotide sequence under stringent conditions. As used herein, "stringent
15 conditions" refers to prewashing in a solution of 6X SSC, 0.2% SDS; hybridizing at 65°C, 6X SSC, 0.2% SDS overnight; followed by two washes of 30 minutes each in 1X SSC, 0.1% SDS at 65°C and two washes of 30 minutes each in 0.2X SSC, 0.1% SDS at 65°C.

20 It is of course expressly contemplated that homologs to the PrAG1 promoter exist in other plants, particularly other coniferous plants, including other members of the *Pinus* genus. Such homologs are also "functionally equivalent variants" of PrAG1 promoter as the phrase is used herein.

25 DNA sequences from plants other than *Pinus radiata* which are homologs of the PrAG1 promoter may be isolated by high throughput sequencing of cDNA libraries prepared from such plants. Alternatively, oligonucleotide probes based on the sequence for the PrAG1 promoter provided in Figure 2 can be synthesized and used to identify positive clones in either cDNA or genomic DNA libraries from other plants
30 by means of hybridization or PCR techniques. Probes should be at least about 10, preferably at least about 15 and most preferably at least about 20 nucleotides in length. Hybridization and PCR techniques suitable for use with such oligonucleotide probes are well known in the art. Positive clones may be analyzed by restriction enzyme digestion, DNA sequencing or the like.

The polynucleotides of the present invention may be generated by synthetic means using techniques well known in the art. Equipment for automated synthesis of oligonucleotides is commercially available from suppliers such as Perkin Elmer/Applied Biosystems Division (Foster City, CA) and may be operated according to the manufacturer's instructions.

The PrAG1 promoter of the invention is plant reproductive-tissue-specific. The primary importance of identification of the polynucleotide of the invention is therefore that it enables the reproductive capacity of plants to be modulated. This modulation will generally involve a reduction in the reproductive capacity of the plant.

Any conventional technique for effecting this can be employed. Examples include co-suppression or anti-sense strategies, a dominant negative approach, or techniques which involve expressing enzymes (such as RNases) to digest, or otherwise be lethal to, RNA post-transcription of a target gene.

Co-suppression can be effected in a manner similar to that discussed, for example, by Napoli *et al* (Plant Cell 2:279-290, 1990) and de Carvalho Niebel *et al* (Plant Cell 7:347-258, 1995). In some cases, it can involve overexpression of the gene of interest through use of multiple constructs of the PrAG1 promoter and gene to be suppressed.

Anti-sense strategies involve using the PrAG1 promoter to effect expression or transcription of DNA with the expression/transcription product being capable of interfering with translation of mRNA transcribed from the gene to be suppressed. This will normally be through the expression/transcription product hybridising to and forming a duplex with the target mRNA.

The expression/transcription product can be a relatively small molecule and still be capable of disrupting mRNA translation. However, the same result is achieved by expressing the target gene in an anti-sense orientation such that the RNA produced

by transcription of the anti-sense oriented gene is complementary to all or part of the endogenous target mRNA.

Such anti-sense strategies are described generally by Robinson-Benion *et al.*,
5 (1995), *Anti-sense techniques, Methods in Enzymol.* 254(23):363-375 and Kawasaki
et al., (1996), *Artific. Organs* 20 (8): 836-848.

Dominant negative approaches involve using the PrAG1 promoter to effect the
expression of a modified DNA binding/activating protein which includes a DNA
10 binding domain but not a activator domain. The result is that the protein binds to
DNA as intended but fails to activate, while at the same time blocking the binding of
the DNA binding/activating peptides which normally bind to the same site.

It is however presently preferred that the reproductive capacity of the plant be
15 reduced or eliminated through the use of the PrAG1 promoter to drive transcription
and expression of a nucleotide sequence which encodes an RNase within the plant
reproductive tissue. Such an approach, in which the PrAG1 promoter is coupled to
the RNase, RNS2, is exemplified herein.

20 To give effect to the above strategies, the invention also provides DNA constructs.
The constructs include the PrAG1 promoter sequence, the DNA intended to be
transcribed/expressed (such as the PrAG1 gene in sense or in anti-sense
orientation or a polynucleotide encoding an RNase) and a termination sequence,
operably linked to the DNA sequence to be transcribed. The promoter sequence is
25 generally positioned at the 5' end of the DNA sequence to be transcribed, and is
employed to initiate transcription of the DNA sequence.

The DNA with which the PrAG1 promoter is operatively associated can encode any
peptide it is desirable to express in plant reproductive tissue. As indicated above,
30 this includes the peptide encoded by PrAG1, but can also be another peptide. That
other peptide can be a peptide which, when produced, causes the reproductive
organs of the plant to abort, redefine themselves as vegetative or stop development.
The peptide encoded can, for example, also be a peptide causing cell death.
Illustrative peptides/genes are Diphtheria Toxin A (DTA), Barnase (from *Bacillus*

amyloliquefaciens), apoptosis genes, glucanase, and RNAses, with the selection of each being a matter of choice for the art skilled worker.

Alternatively, the peptide which is to be expressed under the control of the PrAG1 promoter can be one which, when produced, alters the timing of flowering (ie. either delays or accelerates flowering, such as the ELF-3 and CONSTANS flowering time genes).

The peptide to be expressed can be ligated to the promoter in a sense or antisense orientation, dependant upon the desired effect.

The termination sequence, which is located 3' to the DNA sequence to be transcribed, may come from the same gene as the PrAG1 promoter or may be from a different gene. Many termination sequences known in the art may be usefully employed in the present invention, such as the 3' end of the *Agrobacterium tumefaciens* nopaline synthase gene. However, preferred termination sequences are those from the original gene or from the target species to be transformed.

The DNA constructs of the present invention may also contain a selection marker that is effective in plant cells, to allow for the detection of transformed cells containing the construct. Such markers, which are well known in the art, typically confer resistance to one or more toxins. One example of such a marker is the NPTII gene whose expression results in resistance to kanamycin or hygromycin, antibiotics which are usually toxic to plant cells at a moderate concentration (Rogers *et al.*, in *Methods for Plant Molecular Biology*, A Weissbach and H Weissbach eds, Academic Press Inc., San Diego, CA (1988)). Other examples of markers include visible selection markers such as Green Fluorescent Protein (GFP) and herbicide resistance genes. Alternatively, the presence of the desired construct in transformed cells can be determined without reference to marker genes, by means of other techniques well known in the art, such as Southern and Western blots.

Techniques for operatively linking the components of the DNA constructs are well known in the art and include the use of synthetic linkers containing one or more restriction endonuclease sites as described, for example, by Maniatis *et al.*,

(*Molecular Cloning: A Laboratory Manual*, Cold Spring Harbour Laboratories, Cold Spring Harbour, NY, 1989). The DNA construct may be linked to a vector having at least one replication system, for example, *E. coli*, whereby after each manipulation, the resulting construct can be cloned and sequenced and the correctness of the manipulation determined.

The DNA constructs of the present invention may be used to transform a variety of plants. In one embodiment, these will be plants of the *Pinus* genus. In a preferred embodiment, the DNA constructs are employed to transform *Pinus radiata*, *Pinus taeda*, *Pinus elliotti*, *Pinus clausa*, *Pinus palustris*, *Pinus echinata*, *Pinus ponderosa*, *Pinus jeffrey*, *Pinus resinosa*, *Pinus rigida*, *Pinus banksiana*, *Pinus serotina*, *Pinus strobus*, *Pinus monticola*, *Pinus lambertiana*, *Pinus virginiana*, *Pinus contorta*, *Pinus cariboea*, *Pinus pinaster*, *Pinus brutia*, *Pinus eldarica*, *Pinus coulteri*, *Pinus nigra*, *Pinus sylvestris*, *Pinus tecunumanni*, *Pinus keyisia*, *Pinus oocarpa* and *Pinus maximowii*; and hybrids between the above species.

The constructs can also be used to transform other plants such as trees selected from *Abies amabilis*, *Abies balsamea*, *Abies concolor*, *Abies grandis*, *Abies lasiocarpa*, *Abies magnifica*, *Abies procera*, *Chamaecyparis lawsoniana*, *Chamaecyparis nootkatensis*, *Chamaecyparis thyoides*, *Huniperus virginiana*, *Larix decidua*, *Larix laricina*, *Larix leptolepis*, *Larix occidentalis*, *Larix siberica*, *Libocedrus decurrens*, *Picea abies*, *Picea engelmannii*, *Picea glauca*, *Picea mariana*, *Picea pungens*, *Picea rubens*, *Picea sitchensis*, *Pseudotsuga menziesii*, *Sequoia gigantea*, *Sequoiasempervirens*, *Taxodium distichum*, *Tsuga canadensis*, *Tsuga heterophylla*, *Tsuga mertensiana*, *Thuja occidentalis*, *Thuja plicata*; Eucalypts, such as *Eucalyptus alba*, *Eucalyptus bancroftii*, *Eucalyptus botyroides*, *Eucalyptus bridgesiana*, *Eucalyptus calophylla*, *Eucalyptus camaldulensis*, *Eucalyptus citriodora*, *Eucalyptus cladocalyx*, *Eucalyptus coccifera*, *Eucalyptus curtisii*, *Eucalyptus dalrympleana*, *Eucalyptus deglupta*, *Eucalyptus delagatensis*, *Eucalyptus diversicolor*, *Eucalyptus dunnii*, *Eucalyptus ficifolia*, *Eucalyptus globulus*, *Eucalyptus gomphocephala*, *Eucalyptus gunnii*, *Eucalyptus henryi*, *Eucalyptus laevopinea*, *Eucalyptus macarthurii*, *Eucalyptus macrorhyncha*, *Eucalyptus maculata*, *Eucalyptus marginata*, *Eucalyptus megacarpa*, *Eucalyptus melliodora*, *Eucalyptus nicholii*, *Eucalyptus nitens*, *Eucalyptus nova-anglica*, *Eucalyptus obliqua*, *Eucalyptus obtusiflora*, *Eucalyptus oreades*, *Eucalyptus pauciflora*, *Eucalyptus polybractea*, *Eucalyptus regnans*, *Eucalyptus resinifera*,

Eucalyptus robusta, *Eucalyptus rudis*, *Eucalyptus saligna*, *Eucalyptus sideroxylon*, *Eucalyptus stuartiana*, *Eucalyptus tereticornis*, *Eucalyptus torelliana*, *Eucalyptus urnigera*, *Eucalyptus urophylla*, *Eucalyptus viminalis*, *Eucalyptus viridis*, *Eucalyptus wandoo* and *Eucalyptus youmanni*; and hybrids between any of the above species.

5

As discussed above, transformation of a plant with a DNA construct including an open reading frame coding for a peptide wherein the open reading frame is orientated in a sense direction can, in some cases, lead to a decrease in expression of the peptide by co-suppression. Transformation of the plant with a DNA construct comprising an open reading frame in an anti-sense orientation or a non-coding (untranslated) region of a gene will lead to a decrease in the expression of the peptide in the transformed plant.

Techniques for stably incorporating DNA constructs into the genome of target plants are well known in the art and include *Agrobacterium tumefaciens* mediated introduction, electroporation, protoplast fusion, injection into reproductive organs, injection into immature embryos, high velocity projectile introduction and the like. The choice of technique will depend upon the target plant to be transformed.

Once the cells are transformed, cells having the DNA construct incorporated into their genome may be selected by means of a marker, such as the NPT II or kanamycin resistance markers discussed above. Transgenic cells may then be cultured in an appropriate medium to regenerate whole plants, using techniques well known in the art. In the case of protoplasts, the cell wall is allowed to reform under appropriate osmotic conditions. In the case of seeds or embryos, an appropriate germination or callus initiation medium is employed. For explants, an appropriate regeneration medium is used.

For a review of regeneration of forest trees such as those of the *Pinus* genus, see Dunstan *et al.*, Somatic embryogenesis in woody plants. In: Thorpe, T.A. ed. 1995: *in vitro* embryogenesis of plants. Vol 20 in Current Plant Science and Biotechnology in Agriculture, Chapter 12, pp. 471-540.

The promoter, and constructs containing it, are not restricted in use to plants of forestry species. They can also be used to transform other agronomically important plants in which modulation of reproductive capacity (particularly the timing and abundance of flowering) is desirable. Such plants include cereals, rice, maize, wheat, barley, oats, rye, soyabean and canola.

The resulting transformed plants may be reproduced sexually or asexually, using methods well known in the art, to give successive generations of transgenic plants.

The invention will now be described with reference to the following non-limiting examples.

EXPERIMENTAL

SECTION 1

1. PrAG1 cDNA Cloning:

Total RNA was purified from immature cone of radiata pine according to the protocol of Charles Ainsworth (Plant Molecular Biology Reporter, 12(3), 1994: 198-203). The mRNA was isolated with oligo-T cellulose column. With mRNA as template, the cDNA was synthesised with Cap-Finder cDNA synthesis kit (ClonTech Co.). The cDNA was inserted into Lambda TriplEx Vector (ClonTech Co.), then packaged it with Gigapack@ III packaging extracts (Stratagene Co.) to obtain a cDNA library.

Two degenerate primers were designed:

3' PCR primer: 5' GCIGTIAGIYCITCICCCAT3' ; (SEQ ID NO. 6)

5' PCR primer: 5' AAYCGICARGTIACITT3' (SEQ ID NO. 7)

These primers were used to perform RT-PCR based differential screening on RNA from various female tissue sources, including immature female buds, vegetative tissue from needles and later stages of development. The 50-ul reaction mixture contained 2.5 Units Taq DNA polymerase, 1X Polymerization Buffer (both from

ClonTech Co.), 1mMMgCl₂ , 0.2mMdNTP and 0.25uM primers. The PCR was performed under the following conditions: denaturation at 94°C for 30 s, annealing at 50°C for 1 min and extension at 72°C for 1 min for 30 cycles on a Thermal Cycler 480 (Perkin-Elmer, Norwalk, CT, USA).

5

Fragments were obtained mostly from immature female bud tissue RNA samples. Several DNA fragments were cloned into pGEM-T vector and sequenced. Sequence analysis showed that most of these fragments contained similar sequences. One of the DNA fragments (309 bp) was chosen. This was used as a probe to screen the cDNA library to clone its full-length cDNA and resulted in the cDNA clone, PrAG1. The sequence of PrAG1 was analysed on both strands by the Sanger's dideoxy method (Sanger *et al.*, 1977, Proc. Natl. Acad. Sci. U.S.A. 74: using a Sequenase kit (United States Biochemical co.).

10

15 The resulting sequence is shown in Figure 1 gives the nucleotide sequence coding for the peptide of the invention together with the predicted amino acid sequence.

Sequence comparison and phylogenetic analysis were conducted with the software program MacDNASIS (Version 3.5, Hitachi Corp.). The results of analysis revealed PrAG1 to be a MADS box gene.

20

2. PrAG1 Promoter Cloning:

I. Genomic DNA purification:

25

Genomic DNA was purified from young needles according to a CTAB method as described below.

1) 2 g of young needles of *Pinus radiata* were ground in liquid nitrogen (mortar and pestle) to a fine powder.

30

2) This powder was mixed with 15 mL of pre-warmed CTAB extraction buffer [3% CTAB(W/V), 100mM Tris-HCl pH8.0, 20 mM EDTA pH8.0, 1.4 M NaCl, 1% PVP 940,000, 1% beta mercaptoethanol] and incubated at 65°C for one hour.

3) To the above mixture 15 mL chloroform was added and mixed gently.

4) The contents were centrifuged at 10,000g for 20 minutes at 4°C.

5

5) The supernatant was transferred to a new tube, and mixed with 1/10 volume of 3M sodium acetate (pH4.8), and 0.7 volume of isopropanol. The DNA was precipitated at -20°C for 30 minutes.

10 6) The DNA was pelleted at 10,000 g for 10 minutes at 4°C.

7) The DNA pellet was then air dried and resuspended in 2 mL TE buffer (10 mM Tris-HCl pH7.5, 1mM EDTA pH8.0) and 2 µl of RNase A (10µg/µL) was added. The contents were incubated at 37°C for 30 minutes to remove any RNA from the sample.

15

8) After the incubation, 2 mL of 5M Ammonium acetate and 10 mL of 100% ethanol were added and the contents kept at -20°C for 15 minutes.

20 9) The mixture was then centrifuged at 10,000 g for 10 minutes at 4°C to pellet DNA. The DNA pellet was washed in 70% ethanol twice.

10) The DNA pellet was air dried and resuspended in 200 µL TE buffer.

25 **II. Cloning of PrAG1 Promoter with Two step Genomic DNA Walking.**

1) The Universal Genome Walker Kit (CLONTECH) was employed. For the first step genomic DNA walking, two PrAG1 specific primers were designed and synthesized according to the PrAG1 cDNA sequence. The sequences of the primers were:

30

Primer GSP1: 5' CGC CTT CTT CAA TAA ACC ATT TCG GCG CTT 3' SEQ ID NO. 8

Primer GSP2: 5' GAC CTG TCG GTT CGT AGT ATT TTC AAT CCT 3' SEQ ID NO. 9

2) Based upon the promoter sequence we got from step 1), two PrAG1 promoter sequence specific primer were designed and synthesized. The primers were:

Primer GSP3: 5' TTC GTC CTC CAT TTT GTG CGC TCT CCA TTC 3' SEQ ID NO. 10

5 Primer GSP4: 5' GCA CTC CAC TCT TCC TTT ATT TCT TAC CAC 3'. SEQ ID NO. 11

3) According to the User Manual of Universal Genome Walker Kit, 13 genome walker libraries were constructed after genomic DNA digestion with restriction enzymes: EcoR V, Sca I, Dra I, Pvu II, Ssp I, Stu I, Sma I, Hap I, BsaB I, Bcl136 II, Pml I, Nru I, Hic II.

4) With 13 genome walker libraries as templates, and adaptor primer 1 (AP1 primer from kit) and GSP1 primer, first round PCR was performed under the conditions suggested by the kit manufacturer. After agarose electrophoresis analysis of the PCR product, second round PCR was performed with the nested primers AP2 (Adaptor primer from the kit) and GSP2. The PCR products from the second round PCR were purified and cloned into pGEM-T easy vector (Promega). Following sequence analysis, and DNA sequence comparison with PrAG1 cDNA, one DNA fragment of 1105 bp from Sca I genome walker library was obtained which was identified as the promoter region of PrAG1, based upon the overlapped region between it and PrAG1 cDNA.

5) The second step genome walking was done with primer pair AP1 and GSP3, and primer pair AP2 and GSG4. A DNA fragment of 449 bp from the Dra I genome walker library was identified as the upstream sequence of the PrAG1 promoter cloned from the first step genome walking based on the sequence comparison of overlapped region between them.

6) The 1105bp and 449 bp fragments were used in PCR mediated DNA splicing to synthesize one continuous 1458 bp promoter fragment of PrAG1. This was done as described. One primer was synthesized based on the 5'end sequence of 1105 bp promoter fragment: Primer PLi, 5' AGT TAC TTA ACA ATG CGC AAC CAA GGC 3' (SEQ ID NO. 12). Primer pair PLi and GSP2 was used in PCR to get the promoter fragment of 1105 bp, in which the AP2 primer sequence was removed. This 1105 bp

fragment and 449 bp fragment was then added in one PCR tube as a template with the primer pair of AP2 and GAP2 to do the second round PCR to get the 1458 bp PCR fragment. The conditions of second round PCR were as follows: the first cycle at 95°C for 5 minutes, and 68°C for 10 min; the second cycle at 94°C for 30 seconds (DNA denaturing), DNA annealing at 60 C for 1 min, and DNA synthesis at 72 C for 2 minutes; this regime was cycled 30 times. This 1458 bp fragment was then cloned into pGEM-T easy vector (Promega) and subjected to DNA sequencing on both strands to confirm the DNA sequence and to make sure that no base changes occurred during the PCR process.

The sequence of the promoter, (which is the PrAG1 promoter), is given in Figure 2 from nucleotides 1 to 1320.

7) DNA sequence analysis has indicated that compared to its orthologs from other plants, the PrAG1 showed that there were two possible positions for transcription initiation: at position 791 or 1326 in the Figure 2 sequence. It was found three typical TATA boxes in the PrAG1 promoter at the position of 280 to 286, 282 to 288, 1015 to 1021. Based on the start codon position and short 5' untranslated region in the PrAG1 cDNA, the transcription initiation point is identified as position 1326 in the Figure 2 sequence.

3. DNA and RNA Gel Blot Hybridizations:

Genomic DNA and RNA gel blots were made using standard techniques (Sambrook et al., 1989. Molecular Cloning : A Laboratory Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press).

RNA: Total RNA was prepared from needle, vegetative shoot, stem, immature female cone and immature male cone samples as described above. Briefly, 20µg of total RNA was denatured in formaldehyde loading buffer and fractionated by denaturing agarose gel electrophoresis on a formaldehyde containing gel. The agarose gel was stained with ethidium bromide and a picture taken as control. The RNA was then transferred to a nylon membrane by the capillary blotting method. The RNA was immobilised on the membrane by UV cross-linking and was prehybridized at 65°C

for 2 hours prior to hybridization in 0.5M Na-phosphate, pH 7.2, 7.5%SDS, 1mM EDTA, 100ug/mL salmon sperm DNA. A DNA fragment of PrAG1 3' end region was labelled with ³²P- dCTP (Decaprime II kit, Ambion, Austin, TX), and hybridised to the RNA blot overnight at 65°C. The blot was washed twice in 40mMNa-PO₄, 1%SDS and 1mM EDTA for 30 minutes each at 65°C, and exposed to X-ray film with intensifying screens at -80°C.

DNA: Genomic DNA was prepared from needle tissue with CTAB method. Twenty µg genomic DNA was digested by Bam HI, Bgl II, Eco RI, Hind III and Xba I respectively. After agarose gel running, alkali blotting of DNA to Hybond N membranes was performed as described by the manufacturer (Amersham). The probe hybridisation and washing was as described for the RNA blotting analysis.

The results are shown in Figures 3 and 4.

4. RT-PCR:

Analysis was performed on total RNA isolated from needle, stem, vegetative shoot, immature female cone and immature male cone samples as described above. RNA was reverse-transcribed with MMLV reverse-transcriptase (Gibco BRL) according to the manufacturer's instructions. PCR was performed with two primers: 5'PCR primer (5' TTGTGTACAAATCATGGG3') (SEQ ID NO. 13) and 3'PCR primer (5' GTAAGCCCGTCACCCATC3') (SEQ ID NO. 14). Verification of the specificity of the PCR reactions was achieved through the use of controls that included amplification reaction with single primers, RNase treatment of template, and no template. In those reactions in which no PCR product was detected, the quality of the RNA was tested by UV scanning, and agarose gel electrophoresis. ss-cDNA from the RT reaction was used as a template. The 50-ul reaction mixture contained 2.5 U Taq DNA polymease, 1X Polymerization Buffer (both from ClonTech Co.), 1mM MgCl₂, 0.2mM dNTP and 0.25µM primers. The PCR was performed under following conditions: denaturation at 94°C for 30 s, annealing at 55°C for 1 min and extension at 72°C for 1 min for 30 cycles on Thermal Cycler 480 (Perkin-Elmer, Norwalk, CT, USA). The PCR products were subjected to electrophoresis in agarose gel, and hybridization as described above.

The results are shown in Figure 5.

Discussion

Northern blot hybridization and RT-PCR analysis showed that PrAG1 mRNA is accumulated specifically in the immature female cone and immature male cone; there is no expression detected in needle, stem, and vegetative shoot (Figures 3 and 5). This tissue distribution profile, when combined with the fact that PrAG1 contains a MADS box, verifies that PrAG1 is a reproductive gene in *Pinus radiata* and that the PrAG1 promoter is reproductive-tissue specific.

Southern blot analysis showed that PrAG1 gene exists as a single copy in the genome of *Pinus radiata* (Figure 4).

SECTION 2

Construction of binary vector pRAGPR, plant transformation and regeneration of transgenic tobacco plants

A DNA fragment containing the PrAG1 promoter (1.46 kb, sequence of Figure 2) operably fused to an RNase gene (0.95 kb, RNS2, Taylor *et al. Proc Natl Acad Sci, USA* 90 (11), 5118-5122 (1993)) and containing Hind III and Sac I sites was gel purified and ligated into the Hind III/Sac I sites of binary vector pRD420, containing the NPTII gene for plant selection, (provided by Dr. R.S.S. Datla, PBI, Saskatoon, Canada) resulting in the construct pRAGPR (Figure 6). The construct was introduced into *Agrobacterium tumefaciens* (strain c58 MP90), and used to transform and regenerate *Nicotiana tabacum* var. *Xanthi* by the standard leaf disc transformation method (Horsch *et al.* (1985), A simple and general method for transferring genes into plants. *Science* 227, 1229-1231). Control lines were also generated through leaf disk method without the selection process. After kanamycin selection, the putative transgenic plantlets were rooted in the rooting medium containing kanamycin and then moved to pots containing the Metromix 350 potting mix. Potted plants were maintained under controlled conditions in a growth chamber with 16h photoperiod. The plants were grown through the full life cycle of

the tobacco until senescence and the flowering of the transgenic tobacco assessed relative to controls. Transgenic plants were identified further by PCR with template of genomic DNA and Southern blot analysis to confirm the integration of pRAGPR in transgenic tobacco plants.

5

Polymerase Chain Reaction Amplification

To check the genomic DNA integration of pRAGPR in the transgenic tobacco plants, gene-specific primers for the NPTII gene were employed. The primers used were NPTII-5' primer 5-GAA CAA GAT GGA TTG CAC GC-3' (SEQ ID NO. 15) and NPTII-3' primer 5'-GAA GAA CTC GTC AAG AAG GC-3' (SEQ ID NO. 16). Genomic DNA from each of the control lines and transgenic tobacco lines were isolated from the leaf tissue using the Qiagen DNAeasy kit as per manufacturer's instructions. PCR reactions (50- μ l final volume) were performed using 5 μ l of template DNA. Samples were heated to 95°C for 4 minutes, followed by 35 cycles of 95°C for 45 seconds, 55°C for 30 seconds, and 73°C for 2 minutes, with a final extension step of 73°C for 5 minutes in PTC100 thermal cycler (MJ Research). Amplified DNA fragments were analyzed on a 0.8 agarose gel and visualized by staining with ethidium bromide.

DNA Gel Blot Hybridization

To confirm the genomic integration of PrAG1 promoter-RNase gene cassette in transgenic plants and to determine the copy number, Southern analysis was performed. For Southern analysis, genomic DNA (20 μ g) was digested with appropriate restriction endonucleases, separated by electrophoresis in a 0.8% agarose gel. Two sets of Southern blots were performed, one with digesting the genomic DNA with Hind III and Sac I to drop the PrAG1-RNase cassette and another with Hind III digest alone to test for integration profiles. Following depurination in 0.25 M HCl and denaturation in 0.5 M NaOH, 1.5 M NaCl, the DNA was blotted onto a nylon membrane. The RNase gene-specific probe (which is the whole RNase gene) was radioactively labelled using a random-primed DECAprime II DNA labeling kit (Ambion, Austin, Texas). Filters were hybridized at 65°C in a hybridization buffer containing 0.5 M NaPO₄ (pH 7.5), 7.5% SDS, and 1mM EDTA. All filters were washed finally at 68°C in 20 mM NaPO₄ (pH 7.5), 1% SDS. Filters were then subjected to autoradiography.

Results and Conclusions

Transgenic Tobacco plants expressing PrAG1 promoter controlled RNAs gen .

Eight independent transgenic lines of tobacco (*N. tabaccum* var *Xanthu*) with the PrAG1-RNase fusion construct were obtained together with three control lines processed through the same tissue culture propagation method but without the selection. PCR was done on the putative transgenic plantlets, after rooting in selection medium, to confirm the integration of the NPTII gene. All the eight lines were positive for the NPTII gene and the controls were negative (Figure 7). The transgenic plants were propagated in pots along with the controls.

Southern analysis was done on the eight transgenic lines and the three controls to confirm the integration and profiles of integration of PrAG1promoter-RNase gene cassette. Double digestion with Hind III and Sac I followed by probing with RNase gene probe indicated that all the eight transgenic plants had the PrAG1-RNase cassette (Figure 8). The single digest with Hind III indicated that 5 of the transgenic lines had single integration and three others had two copies of the cassette.

Out of the 5 transgenic plants with single integration two profiles were found based on molecular weight size bands that hybridized to the RNase probe. Three plants had the same size band hybridizing at ~ 4kb (#s 3a, 3 c, 4a) whereas, the two other plants had a band hybridizing at ~3kb range (#s 3b, 4b)(Figure 9).

Of the three transgenic plants that had two copies of the cassette, two had same profile with a band at 5 -6 kb and another at 4 kb hybridizing to the RNase probe (#1a, 1b), whereas the other had two bands, one at 8 kb and another at 4 kb that hybridized to RNase probe (#2) (Figure 9).

All the three controls were negative for hybridization with the RNase probe (Figure 8 and Figure 9).

Reproductive sterility in transgenic tobacco caused by pRAGPR.

All the controls and transgenic lines were allowed to senesce (approximately 3 months). The controls flowered normally whereas none of the transgenic tobacco

containing the pRAGPR construct flowered. The plants with single copy PrAG1-RNase cassette grew at the same rate as controls and senesced at the same time. The transgenic plants with two copies of pRAGPR were slower to grow, and matured late (#1a, 1b and 2).

5

All transgenic plants eventually died and none flowered.

Another phenotypic change observed was in transgenic line #2, which developed additional lateral branches near the top. This may be due to the lack of or decreased apical dominance in these plants. Thus, the inhibition of flowering using the PrAG1-RNase cassette may have an added benefit to increase the biomass of the plant through increased branching if the growing conditions are not limiting.

10

INDUSTRIAL APPLICATION

15

In its primary aspect, the invention provides a new, reproductive-tissue-specific promoter. This promoter can be used in transforming a wide variety of plants. The promoter can also be used to drive expression of any gene which it is desirable to express in plant reproductive organs, including flowering time genes.

20

The invention also has application in modulating, and in particular reducing or eliminating reproductive capacity in plants including those of the *Pinus* genus and *Eucalyptus* genus. Such plants have utility in forestry.

25

The availability of reproductively null or sterile pine or eucalyptus trees has the additional advantage that it will be possible to introduce further exogenous genetic material into those trees without the risk that the material will be passed on to other trees.

30

Those persons skilled in the art will appreciate that the specific description provided is exemplary only, and that modifications and variations may be made without departing from the scope of the invention.

CLAIMS

1. A polynucleotide which has the nucleotide sequence of SEQ ID NO. 1 and which has the ability, when operatively associated with a nucleotide sequence encoding a peptide, to promote transcription of that nucleotide sequence, or a polynucleotide which is a functionally equivalent variant thereof.
2. A plant reproductive tissue promoter which has the nucleotide sequence of SEQ ID NO. 1 or a functionally equivalent variant thereof.
3. A plant reproductive tissue promoter which has the nucleotide sequence of SEQ ID NO. 2.
4. A DNA construct which comprises:
- (a) a polynucleotide having activity as a transcriptional promoter according to claim 1;
 - (b) an open reading frame polynucleotide coding for a peptide; and
 - (c) a termination sequence.
5. A DNA construct which comprises:
- (a) a promoter sequence according to claim 2 or claim 3;
 - (b) an open reading frame polynucleotide coding for a peptide; and
 - (c) a termination sequence.
6. A construct as claimed in claim 4 or claim 5 in which the open reading frame is in a sense orientation.
7. A construct according to claim 4 or claim 5 in which the open reading frame is an anti-sense orientation.
8. A construct according to any one of claims 4-7 wherein said open reading frame polynucleotide encodes a peptide having SEQ ID NO. 3.

9. A construct according to any one of claims 4-7 wherein said open reading frame polynucleotide encodes a peptide which, when expressed in reproductive tissue of a plant, causes said plant's reproductive organs to abort.

5

10. A construct according to any one of claims 4-7 wherein said open reading frame polynucleotide encodes a peptide which, when expressed in reproductive tissue of a plant, causes said plant's reproductive organs to redefine themselves as vegetative.

10

11. A construct according to any one of claims 4-7 wherein said open reading frame polynucleotide encodes a peptide which, when expressed in reproductive tissue of a plant, causes said plant's reproductive organs to stop development.

15

12. A construct according to any one of claims 4-7 wherein said open reading frame polynucleotide encodes a peptide which, when expressed in reproductive tissue of a plant, causes cell death.

20

13. A construct according to claim 12 wherein the peptide which causes cell death is selected from diphtheria toxin A and Barnase.

14. A construct according to claim 12 wherein the peptide which causes cell death is an RNase.

25

15. A construct according to claim 14 wherein said RNase is encoded by the nucleotide sequence of SEQ ID NO. 5.

30

16. A construct according to any one of claims 4-7 wherein said open reading frame polynucleotide encodes a peptide, which when expressed in reproductive tissue of a flowering plant, causes an alteration in the timing of flowering of said plant.

17. A construct according to any one of claims 4-16 which further includes:

(d) a selection marker sequence.

18. A construct according to claim 17 in which said selection marker sequence is the NPTII gene.

5

19. A transgenic plant cell which includes a construct according to any one of claims 4-18.

10

20. A transgenic plant which includes a construct according to any one of claims 4-18.

21. A transgenic plant which contains a polynucleotide according to claim 1 or a promoter according to claim 2 or claim 3, which plant has a reduced reproductive capacity.

15

22. A transgenic plant according to claim 21 wherein in said plant said polynucleotide or promoter is operatively associated with a nucleotide sequence encoding a peptide, which when expressed in reproductive tissue of the plant, causes the plant's reproductive organs to abort, redefine as vegetative or stop development.

20

23. A transgenic plant according to claim 21 wherein in said plant said polynucleotide or promoter is operatively associated with a nucleotide sequence encoding a RNase.

25

24. A transgenic plant according to claim 23 in which the RNase has the sequence of SEQ ID NO. 5.

30

25. A transgenic plant according to any one of claims 20-24 wherein said plant is a coniferous plant.

26. A transgenic plant according to claim 25 which is a coniferous plant of the *Pinus* genus.

27. A transgenic plant according to claim 26 which is a member of a species selected from *Pinus radiata*, *Pinus taeda*, *Pinus elliotti*, *Pinus clausa*, *Pinus palustris*, *Pinus echinata*, *Pinus ponderosa*, *Pinus jeffrey*, *Pinus resinosa*, *Pinus rigida*, *Pinus banksiana*, *Pinus serotina*, *Pinus strobus*, *Pinus monticola*, *Pinus lambertiana*, *Pinus virginiana*, *Pinus contorta*, *Pinus cariboea*, *Pinus pinaster*, *Pinus brutia*, *Pinus eldarica*, *Pinus coulteri*, *Pinus nigra*, *Pinus sylvestris*, *Pinus tecunumannii*, *Pinus keyisia*, *Pinus oocarpa* and *Pinus maximowii*; and hybrids between any of the above species.

28. A transgenic plant according to any one of claims 20-24 which is a tree.

29. A transgenic plant according to claim 28 which is a member of the *Eucalyptus* genus.

30. A transgenic plant according to claim 29 which is a member of a species selected from; *Eucalyptus alba*, *Eucalyptus bancroftii*, *Eucalyptus botyroides*, *Eucalyptus bridgesiana*, *Eucalyptus calophylla*, *Eucalyptus camaldulensis*, *Eucalyptus citriodora*, *Eucalyptus cladocalyx*, *Eucalyptus coccifera*, *Eucalyptus curtisii*, *Eucalyptus dalrympleana*, *Eucalyptus deglupta*, *Eucalyptus delagatensis*, *Eucalyptus diversicolor*, *Eucalyptus dunnei*, *Eucalyptus ficifolia*, *Eucalyptus globulus*, *Eucalyptus gomphocephala*, *Eucalyptus gunnii*, *Eucalyptus henryi*, *Eucalyptus laevopinea*, *Eucalyptus macarthurii*, *Eucalyptus macrorhyncha*, *Eucalyptus maculata*, *Eucalyptus marginata*, *Eucalyptus megacarpa*, *Eucalyptus melliodora*, *Eucalyptus nicholii*, *Eucalyptus nitens*, *Eucalyptus nova-anglica*, *Eucalyptus obliqua*, *Eucalyptus obtusiflora*, *Eucalyptus oreades*, *Eucalyptus pauciflora*, *Eucalyptus polybractea*, *Eucalyptus regnans*, *Eucalyptus resinifera*, *Eucalyptus robusta*, *Eucalyptus rudis*, *Eucalyptus saligna*, *Eucalyptus sideroxylon*, *Eucalyptus stuartiana*, *Eucalyptus tereticornis*, *Eucalyptus torelliana*, *Eucalyptus urnigera*, *Eucalyptus urophylla*, *Eucalyptus viminalis*, *Eucalyptus viridis*, *Eucalyptus wandoo* and *Eucalyptus youmanni*; and hybrids between any of the above species.

1/7

TGT	GTA	CAA	ATC	ATG	GGT	CGT	GGG	AAG	ATT	GAG	ATA	AAG	AGG	ATT	GAA	AAT	ACT	54
				M	G	R	G	K	I	E	I	K	R	I	E	N	T	
ACG	AAC	CGA	CAG	GTC	ACT	TTC	TGC	AAG	CGC	CGA	AAT	GGT	TTA	TTA	AAG	AAG	GCG	108
T	N	R	Q	V	T	F	C	K	R	R	N	G	L	L	K	K	A	
TAT	GAA	TTA	TCA	GTT	CTT	TGT	GAT	GCA	GAA	GTG	GCC	CTC	ATC	GTC	TTC	TCC	AGC	162
Y	E	L	S	V	L	C	D	A	E	V	A	L	I	V	F	S	S	
AGA	GGG	AGA	CTT	TAT	GAA	TTT	GCC	AAC	CAC	AGC	GTG	AAG	AGG	ACG	ATT	GAG	AGG	216
R	G	R	L	Y	E	F	A	N	H	S	V	K	R	T	I	E	R	
TAC	AAG	AAG	ACT	TGC	GTT	GAC	AAC	AAC	CAC	GGA	GGG	GCG	ATA	TCA	GAG	TCC	AAT	270
Y	K	K	T	C	V	D	N	N	H	G	G	A	I	S	E	S	N	
TCT	CAG	TAT	TGG	CAA	CAG	GAG	GCT	GGT	AAA	CTC	AGA	CAA	CAG	ATT	GAC	ATT	TTG	324
S	Q	Y	W	Q	Q	E	A	G	K	L	R	Q	Q	I	D	I	L	
CAA	AAT	GCA	AAT	AGG	CAT	TTG	ATG	GGT	GAC	GGG	CTT	ACA	GCT	TTG	AAC	ATT	AAG	378
Q	N	A	N	R	H	L	M	G	D	G	L	T	A	L	N	I	K	
GAA	CTC	AAG	CAA	CTT	GAG	GTT	CGA	CTT	GAA	AAA	GGA	ATC	AGC	CGA	GTG	CGA	TCC	432
E	L	K	Q	L	E	V	R	L	E	K	G	I	S	R	V	R	S	
AAA	AAG	AAC	GAG	ATG	TTG	CTT	GAA	GAG	ATC	GAC	ATC	ATG	CAG	AGA	AGG	GAA	CAC	486
K	K	N	E	M	L	L	E	E	I	D	I	M	Q	R	R	E	H	
ATA	CTT	ATC	CAG	GAG	AAT	GAG	ATT	CTT	CGC	AGC	AAG	ATA	GCC	GAG	TGT	CAG	AAT	540
I	L	I	Q	E	N	E	I	L	R	S	K	I	A	E	C	Q	N	
AGC	CAC	AAC	ACG	AAC	ATG	TTA	TCA	GCT	CCG	GAA	TAT	GAT	GCA	CTG	CCC	GCA	TTC	594
S	H	N	T	N	M	L	S	A	P	E	Y	D	A	L	P	A	F	
GAC	TCT	CGA	AAT	TTC	CTA	CAT	GCA	AAT	CTA	ATC	GAT	GCG	GCC	CAT	CAC	TAT	GCA	648
D	S	R	N	F	L	H	A	N	L	I	D	A	A	H	H	Y	A	
CAT	CAG	GAA	CAA	ACA	ACG	CTT	CAG	CTT	GGC	TGA	ACG	TTG	AAG	CGG	TGG	ACG	CTT	702
H	Q	E	Q	T	T	L	Q	L	G	*								
AAA	ACT	CAA	TCA	AGG	CAC	CCG	AAA	AAT	ATG	CTA	GTA	ACC	TTG	AAT	GAG	ATT	CAG	756
AGT	CGA	AAT	ATT	GCG	AGG	CAA	GAG	CAC	AAT	GGA	AGA	GAT	AGC	TCC	TAG	TAT	GAA	810
TAT	GGA	TTT	ATG	ATA	TTA	ACA	TAT	GGT	TTG	TCA	GCT	TTA	AAT	ATA	GCT	GTT	TGA	864
AAC	AAA	GAA	TAC	AAC	ATA	TTA	GCT	AGT	ATT	TTT	TTG	GCG	CAT	GTT	ATC	TTT	CTG	918
TTG																		921

FIGURE 1

2/7

AAACTCGACAGCAAATATGATTTAGATTATGACCTAGAAATAAGCATAGCATTAAAGCAT 60
 ATACATAACAAGCGGTGATATACTCTGACTGCCACTGTACTTGAGGAAAGGTAGTGGACT 120
 CTGCTCAGGTACATTAGTTTGGTAAGGTTGGCTTGGCTTCTGGGTAATATGAGAAGTAAA 180
 GAAGTAAAAGGTATTTGACTCTAGTCAAGTACATTGGATTGCCTTTGTCTGGGGCTTGGAT 240
 GGCTTGGGTTTCGTGTGAGAAGCCAACAATTTATAAGAAATATATAAAATAAAAAATAAAA 300
 AAATTTAAGTGTTGGAAGTGAAAACGGTGGGGCAGAAATATACACAGAAGAGTACTTTAA 360
 CAATGCGCAACCAAGGCAGATTCACAACCTTGATTTCTGGACCTCGAATACGAGATAATGG 420
 TGGTAAGAAATAAAGGAAGAGTGGAGTGCATTTGAAAATGAATGGAGAGCGCACAAAATG 480
 GAGGACGAATAAAATGAAATATAATGCAAGAGTGCATTTCCCTATTATTTCCAGAAATGTA 540
 TATGTGGGGTCTGGCATTACATGGGCGTCGCATTCAGGGGGTGTCTAGCGGTCTTTGA 600
 TTGCAGTGTGGGAGTTGCAACATGTACCAACAAATCCATTCATCCCAAACCTAAATTTA 660
 TCCTCTCCATTACTATTACCTACACCTATACCTAGTAAATATGTCCTGCCTTGTAACCTCC 720
 TCCACTGCCTGCACACGTCTTAGTCAATCCATCTGCCTTCAAATAGGCATTATTTTGTTC 780
 TTTCCCCTCCGACTGAAAGGCTATCGACCGACCGACCGCTCATCTTCTTCTTCTGCGCAA 840
 TTTTTTCTGCTGGATCATCATATTACCATCATCGCCATCCCCACCATCATCATCATGAT 900
 GGTATCTCTATCTCTCCCTGGCAATCGATTGTAGAGGAAAGGAAGAGGGAAGGGGCATAT 960
 GTATTGATCAACCTACCCGAAAAACAATCTGATCAGCCCTGCTCAATCTTGCTTATAAA 1020
 TCTCTTATCCACTGTTCAATCATTCAGGTTTCTTCCCACTTTCAAGCAAAGGCGCCCGGA 1080
 TTGGCCGTGTTCTTAGATTTTCAGGTACTTAAATGGACAATATTCCCCACCTGAAGCCGT 1140
 TCTGAAAAAGATTTGTTTGTAGAAACAAACGATTGTAATATTTGCTTAAGTTGAGCTTAA 1200
 GGGGTTTGGTACCTAACTTGCCTTGTGGTTATTTGTTTCTCAGAACTCGGGCTGCGTCCA 1260
 ACTGTAGGAACGAACCAGCACAAGGGGTTGCAGCTTTTGCTGTTGCTGTTGCGCCCATTG 1320
 CTTTTGGACTGGTATTAGTAGTTGCAGCTTTGTTTTGCATACGCTGTGAGGATCTGTGCG 1380
 CGGAAATTTTGTGTACAAATC 1401

FIGURE 2




FIGURE 3




FIGURE 4

5/7




FIGURE 5

6/7




FIGURE 6

7/7

Figure 7



Figure 8





Figure 9



SEQUENCE LISTING

<110> Carter Holt Harvey Limited
 Tasman Biotechnology Limited
 Michigan Technological University
 Podila, Gopi Krishna
 Liu, Jun-Jun
 Karnosky, David F

<120> Plants Having Modified Reproductive Activity

<130> 25635 MRB

<140>

<141>

<150> NZ334715

<151> 1999-03-17

<160> 17

<170> PatentIn Ver. 2.1

<210> 1

<211> 1320

<212> DNA

<213> Pinus radiata

<400> 1

```

aaactcgaca gcaaatatga tttagattat gacctagaaa taagcatagc attaaagcat 60
atacataaca agcggtgata tactctgact gccactgtac ttgaggaaag gtagtggact 120
ctgctcaggt acattagttt ggtaagggtg gcttggcttc tgggtaatat gagaagtaaa 180
gaagtaaaaag gtatttgact ctagtcaagt acattggatt gcctttgtcg gggcttggat 240
ggcttgggtt cgtgtgagaa gccaacaatt tataagaaat atataaaata aaaaataaaa 300
aaatttaagt gttggaagtg aaaacggtgg ggcagaaata tacacagaag agtacttta 360
caatgcgcaa ccaaggcaga ttcacaactt gatttctgga cctcgaatac gagataatgg 420
tggtaagaaa taaaggaaga gtggagtgc tttgaaaatg aatggagagc gcacaaaatg 480
gaggacgaat aaatgaaata taatgcaaga gtgcatttcc ctattatttc cagaaatgta 540
tatgtggggt cggcattcac atgggcgtcg cattcagggg gtgtcatagc ggtcctttga 600
ttgcagtgtg ggagttgcaa catgtaccaa caaatccatt catcccaaaa cctaaattta 660
tcctctccat tactattacc tacacctata cctagtaaat atgtcctgcc ttgtaactcc 720
tccactgcct gcacacgtct tagtcaatcc atctgccttc aaataggcat tattttgttc 780
tttcccctcc gactgaaagg ctatcgaccg accgaccgct catcttcttc ttctgcgcaa 840
ttttttctgc tggatcatca tcattaccat catcgccatc cccaccatca tcatcatgat 900
ggtatctcta tctctccctg gcaatcgatt gtagaggaaa ggaagaggga aggggcatat 960
gtattgatca acctaccga aaaaacaatc tgatcagccc tgctcaatct tgcttataaa 1020
tctcttatcc actgttcaat cattcaggtt tcttcccact ttcaagcaaa ggcgccccga 1080
ttggccgtgt tcttagattt tcaggtactt aaatggacaa tattccccac ctgaagccgt 1140

```

```
tctgaaaaag atttgtttgt agaaacaaac gattgtaata tttgcttaag ttgagcttaa 1200
gggggtttggt acctaacttg ccttggtggtt atttgtttct cagaactcgg gctgcgtcca 1260
actgtaggaa cgaaccagca caaggggttg cagcttttgc tgttgctgtt gcgcccattg 1320
```

<210> 2

<211> 1401

<212> DNA

<213> Pinus radiata

<400> 2

```
aaactcgaca gcaaatatga tttagattat gacctagaaa taagcatagc attaaagcat 60
atacataaca agcgggtgata tactctgact gccactgtac ttgaggaaaag gtagtggact 120
ctgctcaggt acattagttt ggtaagggtt gcttggcttc tgggtaatat gagaagtaaa 180
gaagtaaaag gtatttgact ctagtcaagt acattggatt gcctttgtcg gggcttggat 240
ggcttgggtt cgtgtgagaa gccacaatt tataagaaat atataaaata aaaaataaaa 300
aaatttaagt gttggaagtg aaaacgggtg ggcagaaata tacacagaag agtacttta 360
caatgcgcaa ccaaggcaga ttcacaactt gatttctgga cctcgaatac gagataatgg 420
tggtaaagaa taaaggaaga gtggagtgc tttgaaaatg aatggagagc gcacaaaatg 480
gaggacgaat aaatgaaata taatgcaaga gtgcatttcc ctattatttc cagaaatgta 540
tatgtggggt cggcattcac atgggcgtcg cattcagggg gtgtcatagc ggtcctttga 600
ttgcagtgtg ggagttgcaa catgtacca caaatccatt catcccaaaa cctaaattta 660
tcctctccat tactattacc tacacctata cctagtaa atgtcctgcc ttgtaactcc 720
tccactgcct gcacacgtct tagtcaatcc atctgccttc aaataggcat tattttgttc 780
tttccccctc gactgaaagg ctatcgaccg accgaccgct catcttcttc ttctgcgcaa 840
ttttttctgc tggatcatca tcattaccat catcgccatc cccaccatca tcatcatgat 900
ggtatctcta tctctccctg gcaatcgatt gtagaggaaa ggaagagggg aggggcatat 960
gtattgatca acctaccgga aaaaacaatc tgatcagccc tgctcaatct tgcttataaa 1020
tctcttatcc actgttcaat cattcaggtt tcttcccact ttcaagcaaa ggcgccccga 1080
ttggccgtgt tcttagattt tcaggtaactt aaatggacaa tattccccac ctgaagccgt 1140
tctgaaaaag atttgtttgt agaaacaaac gattgtaata tttgcttaag ttgagcttaa 1200
gggggtttggt acctaacttg ccttggtggtt atttgtttct cagaactcgg gctgcgtcca 1260
actgtaggaa cgaaccagca caaggggttg cagcttttgc tgttgctgtt gcgcccattg 1320
cttttggact ggtattagta gttgcagctt tgttttgc atcgctgtgag gatctgtgcg 1380
cggaaatttt gtgtacaaat c 1401
```

<210> 3

<211> 909

<212> DNA

<213> Pinus radiata

<220>

<221> CDS

<222> (1)..(669)

<223> Pinus radiata MADS box protein mRNA, complete cds

<300>

<301> Jun-Jun, Liu
 Podila, G K.
 <302> Not applicable
 <303> Direct submission
 <304> -
 <305> -
 <306> ---
 <307> 1997-09-09
 <308> Genbank AF023615
 <309> 1999-01-26
 <313> 1 TO 909

<400> 3

atg ggt cgt ggg aag att gag ata aag agg att gaa aat act acg aac	48
Met Gly Arg Gly Lys Ile Glu Ile Lys Arg Ile Glu Asn Thr Thr Asn	
1 5 10 15	
 cga cag gtc act ttc tgc aag cgc cga aat ggt tta tta aag aag gcg	96
Arg Gln Val Thr Phe Cys Lys Arg Arg Asn Gly Leu Leu Lys Lys Ala	
20 25 30	
 tat gaa tta tca gtt ctt tgt gat gca gaa gtg gcc ctc atc gtc ttc	144
Tyr Glu Leu Ser Val Leu Cys Asp Ala Glu Val Ala Leu Ile Val Phe	
35 40 45	
 tcc agc aga ggg aga ctt tat gaa ttt gcc aac cac agc gtg aag agg	192
Ser Ser Arg Gly Arg Leu Tyr Glu Phe Ala Asn His Ser Val Lys Arg	
50 55 60	
 acg att gag agg tac aag aag act tgc gtt gac aac aac cac gga ggg	240
Thr Ile Glu Arg Tyr Lys Lys Thr Cys Val Asp Asn Asn His Gly Gly	
65 70 75 80	
 gcg ata tca gag tcc aat tct cag tat tgg caa cag gag gct ggt aaa	288
Ala Ile Ser Glu Ser Asn Ser Gln Tyr Trp Gln Gln Glu Ala Gly Lys	
85 90 95	
 ctc aga caa cag att gac att ttg caa aat gca aat agg cat ttg atg	336
Leu Arg Gln Gln Ile Asp Ile Leu Gln Asn Ala Asn Arg His Leu Met	
100 105 110	
 ggt gac ggg ctt aca gct ttg aac att aag gaa ctc aag caa ctt gag	384
Gly Asp Gly Leu Thr Ala Leu Asn Ile Lys Glu Leu Lys Gln Leu Glu	
115 120 125	
 gtt cga ctt gaa aaa gga atc agc cga gtg cga tcc aaa aag aac gag	432
Val Arg Leu Glu Lys Gly Ile Ser Arg Val Arg Ser Lys Lys Asn Glu	
130 135 140	

atg ttg ctt gaa gag atc gac atc atg cag aga agg gaa cac ata ctt 480
 Met Leu Leu Glu Glu Ile Asp Ile Met Gln Arg Arg Glu His Ile Leu
 145 150 155 160

atc cag gag aat gag att ctt cgc agc aag ata gcc gag tgt cag aat 528
 Ile Gln Glu Asn Glu Ile Leu Arg Ser Lys Ile Ala Glu Cys Gln Asn
 165 170 175

agc cac aac acg aac atg tta tca gct ccg gaa tat gat gca ctg ccc 576
 Ser His Asn Thr Asn Met Leu Ser Ala Pro Glu Tyr Asp Ala Leu Pro
 180 185 190

gca ttc gac tct cga aat ttc cta cat gca aat cta atc gat gcg gcc 624
 Ala Phe Asp Ser Arg Asn Phe Leu His Ala Asn Leu Ile Asp Ala Ala
 195 200 205

cat cac tat gca cat cag gaa caa aca acg ctt cag ctt ggc tga 669
 His His Tyr Ala His Gln Glu Gln Thr Thr Leu Gln Leu Gly
 210 215 220

acgttgaagc ggtggacgct taaaactcaa tcaaggcacc cgaaaaatat gctagtaacc 729

ttgaatgaga ttcagagtcg aaatattgcg aggcaagagc acaatggaag agatagctcc 789

tagtatgaat atggatttat gatattaaca tatggtttgt cagctttaaa tatagctgtt 849

tgaaacaaag aatacaacat attagctagt attttttttg gcgatgttat ctttctgttg 909

<210> 4

<211> 222

<212> PRT

<213> Pinus radiata

<400> 4

Met Gly Arg Gly Lys Ile Glu Ile Lys Arg Ile Glu Asn Thr Thr Asn
 1 5 10 15
 Arg Gln Val Thr Phe Cys Lys Arg Arg Asn Gly Leu Leu Lys Lys Ala
 20 25 30
 Tyr Glu Leu Ser Val Leu Cys Asp Ala Glu Val Ala Leu Ile Val Phe
 35 40 45
 Ser Ser Arg Gly Arg Leu Tyr Glu Phe Ala Asn His Ser Val Lys Arg
 50 55 60
 Thr Ile Glu Arg Tyr Lys Lys Thr Cys Val Asp Asn Asn His Gly Gly
 65 70 75 80
 Ala Ile Ser Glu Ser Asn Ser Gln Tyr Trp Gln Gln Glu Ala Gly Lys
 85 90 95

```

Leu Arg Gln Gln Ile Asp Ile Leu Gln Asn Ala Asn Arg His Leu Met
      100                      105                      110
Gly Asp Gly Leu Thr Ala Leu Asn Ile Lys Glu Leu Lys Gln Leu Glu
      115                      120                      125
Val Arg Leu Glu Lys Gly Ile Ser Arg Val Arg Ser Lys Lys Asn Glu
      130                      135                      140
Met Leu Leu Glu Glu Ile Asp Ile Met Gln Arg Arg Glu His Ile Leu
145                      150                      155                      160
Ile Gln Glu Asn Glu Ile Leu Arg Ser Lys Ile Ala Glu Cys Gln Asn
      165                      170                      175
Ser His Asn Thr Asn Met Leu Ser Ala Pro Glu Tyr Asp Ala Leu Pro
      180                      185                      190
Ala Phe Asp Ser Arg Asn Phe Leu His Ala Asn Leu Ile Asp Ala Ala
      195                      200                      205
His His Tyr Ala His Gln Glu Gln Thr Thr Leu Gln Leu Gly
      210                      215                      220

```

<210> 5

<211> 1012

<212> DNA

<213> Arabidopsis thaliana

<220>

<221> CDS

<222> (16)..(795)

<223> Arabidopsis thaliana ribonuclease (RNS2) mRNA,
complete cds

<300>

<301> Taylor, C B.

Bariola, P A.

delCardayre, S B.

Raines, R T.

Green, P J.

<302> RNS2: a senescence-associated RNase of Arabidopsis that
diverged from the S-RNases before speciation

<303> Proc. Natl. Acad. Sci. U.S.A.

<304> 90

<305> 11

<306> 5118-5122

<307> 1993

<308> Genbank

<309> 1994-10-30

<313> 1 TO 1012

<400> 5

atcgaattaa agtca atg gcg tca cgt tta tgt ctt ctc ctt ctc gtt gcg 51
 Met Ala Ser Arg Leu Cys Leu Leu Leu Leu Val Ala
 1 5 10

tgt atc gcc gga gca ttt gcc gga gac gtc atc gaa ctc aat cga tct 99
 Cys Ile Ala Gly Ala Phe Ala Gly Asp Val Ile Glu Leu Asn Arg Ser
 15 20 25

cag agg gag ttc gat tat ttc gct cta tct ctt caa tgg cct gga acc 147
 Gln Arg Glu Phe Asp Tyr Phe Ala Leu Ser Leu Gln Trp Pro Gly Thr
 30 35 40

tat tgc cgt gga act cgc cat tgt tgc tcc aaa aac gct tgc tgc aga 195
 Tyr Cys Arg Gly Thr Arg His Cys Cys Ser Lys Asn Ala Cys Cys Arg
 45 50 55 60

ggc tcc gat gct cca act caa ttc aca att cat ggg tta tgg cct gac 243
 Gly Ser Asp Ala Pro Thr Gln Phe Thr Ile His Gly Leu Trp Pro Asp
 65 70 75

tat aac gat ggt tgc tgg cct tca tgt tgt tat cga tct gac ttt aaa 291
 Tyr Asn Asp Gly Ser Trp Pro Ser Cys Cys Tyr Arg Ser Asp Phe Lys
 80 85 90

gag aag gag att tca acg ttg atg gat ggt ctt gag aag tac tgg cct 339
 Glu Lys Glu Ile Ser Thr Leu Met Asp Gly Leu Glu Lys Tyr Trp Pro
 95 100 105

agt ctc agt tgt ggt tct cca tca tca tgc aat ggt ggg aaa ggg tca 387
 Ser Leu Ser Cys Gly Ser Pro Ser Ser Cys Asn Gly Gly Lys Gly Ser
 110 115 120

ttt tgg ggc cac gag tgg gag aaa cat ggg act tgt tct tct cct gtt 435
 Phe Trp Gly His Glu Trp Glu Lys His Gly Thr Cys Ser Ser Pro Val
 125 130 135 140

ttt cat gat gag tat aat tac ttc ctt acc aca ctt aat ctc tac ttg 483
 Phe His Asp Glu Tyr Asn Tyr Phe Leu Thr Thr Leu Asn Leu Tyr Leu
 145 150 155

aag cat aat gtc acg gat gtc ctt tat caa gct ggc tat gtt gct tcc 531
 Lys His Asn Val Thr Asp Val Leu Tyr Gln Ala Gly Tyr Val Ala Ser
 160 165 170

aac agt gaa aag tat cct cta gga ggt atc gta aca gcc att cag aat 579
 Asn Ser Glu Lys Tyr Pro Leu Gly Gly Ile Val Thr Ala Ile Gln Asn
 175 180 185

```

gca ttt cat atc acc cct gaa gtg gtt tgc aaa aga gat gca atc gat 627
Ala Phe His Ile Thr Pro Glu Val Val Cys Lys Arg Asp Ala Ile Asp
    190                      195                      200

gaa ata cgt ata tgc ttc tat aaa gat ttt aag ccc agg gac tgt gtt 675
Glu Ile Arg Ile Cys Phe Tyr Lys Asp Phe Lys Pro Arg Asp Cys Val
    205                      210                      215                      220

ggg tca caa gat ttg aca tct aga aag tca tgc ccc aag tac gta agt 723
Gly Ser Gln Asp Leu Thr Ser Arg Lys Ser Cys Pro Lys Tyr Val Ser
                      225                      230                      235

ttg ccg gaa tac acg cca tta gat ggt gaa gct atg gtt ctg aag atg 771
Leu Pro Glu Tyr Thr Pro Leu Asp Gly Glu Ala Met Val Leu Lys Met
                      240                      245                      250

cca aca gaa aga gaa gct ctt tga atcggaaaag atgggagctt tgttatcttc 825
Pro Thr Glu Arg Glu Ala Leu
    255                      260

tgagagacaa tacatacatg tctctgatgt tgtaacttta ctaccaaaac ctataaagat 885

tggcttattt cgttctattg gatatgtatc atcattactg gtaaatacaag tttctttcta 945

ataatgtaga agatcagaaa atccataaga agatatcaac atttgagttc tatggtaaaaa 1005

aaaaaaaaa 1012

```

```

<210> 6
<211> 259
<212> PRT
<213> Arabidopsis thaliana

```

```

<400> 6
Met Ala Ser Arg Leu Cys Leu Leu Leu Leu Val Ala Cys Ile Ala Gly
  1           5           10           15
Ala Phe Ala Gly Asp Val Ile Glu Leu Asn Arg Ser Gln Arg Glu Phe
          20           25           30
Asp Tyr Phe Ala Leu Ser Leu Gln Trp Pro Gly Thr Tyr Cys Arg Gly
          35           40           45
Thr Arg His Cys Cys Ser Lys Asn Ala Cys Cys Arg Gly Ser Asp Ala
          50           55           60
Pro Thr Gln Phe Thr Ile His Gly Leu Trp Pro Asp Tyr Asn Asp Gly
          65           70           75           80
Ser Trp Pro Ser Cys Cys Tyr Arg Ser Asp Phe Lys Glu Lys Glu Ile
          85           90           95
Ser Thr Leu Met Asp Gly Leu Glu Lys Tyr Trp Pro Ser Leu Ser Cys

```

100							105					110				
Gly	Ser	Pro	Ser	Ser	Cys	Asn	Gly	Gly	Lys	Gly	Ser	Phe	Trp	Gly	His	
115							120					125				
Glu	Trp	Glu	Lys	His	Gly	Thr	Cys	Ser	Ser	Pro	Val	Phe	His	Asp	Glu	
130							135					140				
Tyr	Asn	Tyr	Phe	Leu	Thr	Thr	Leu	Asn	Leu	Tyr	Leu	Lys	His	Asn	Val	
145							150					155				
Thr	Asp	Val	Leu	Tyr	Gln	Ala	Gly	Tyr	Val	Ala	Ser	Asn	Ser	Glu	Lys	
165							170					175				
Tyr	Pro	Leu	Gly	Gly	Ile	Val	Thr	Ala	Ile	Gln	Asn	Ala	Phe	His	Ile	
180							185					190				
Thr	Pro	Glu	Val	Val	Cys	Lys	Arg	Asp	Ala	Ile	Asp	Glu	Ile	Arg	Ile	
195							200					205				
Cys	Phe	Tyr	Lys	Asp	Phe	Lys	Pro	Arg	Asp	Cys	Val	Gly	Ser	Gln	Asp	
210							215					220				
Leu	Thr	Ser	Arg	Lys	Ser	Cys	Pro	Lys	Tyr	Val	Ser	Leu	Pro	Glu	Tyr	
225							230					235				
Thr	Pro	Leu	Asp	Gly	Glu	Ala	Met	Val	Leu	Lys	Met	Pro	Thr	Glu	Arg	
245							250					255				
Glu	Ala	Leu														

<210> 7

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Made in lab

<220>

```
<221> modified_base
```

<222> (3)

<223> i

<220>

<221> modified_base

<222> (6)

<223> i

<220>

<221> modified_base

<222> (9)

<223> i

<220>

```
<221> modified_base
```


<222> (12)

<223> i

<220>

<221> modified_base

<222> (15)

<223> i

<400> 7

gcngtnagny cntcncccat

20

<210> 8

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Made in lab

<220>

<221> modified_base

<222> (6)

<223> i

<220>

<221> modified_base

<222> (12)

<223> i

<220>

<221> modified_base

<222> (15)

<223> i

<400> 8

aaycgncarg tnacntt

17

<210> 9

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Made in lab

<400> 9

cgctttcttc aataaaccat ttcggcgctt

30

<210> 10

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Made in lab

<400> 10

gacctgtcgg ttcgtagtat tttcaatcct

30

<210> 11

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Made in lab

<400> 11

ttcgtcctcc attttgtgcg ctctccattc

30

<210> 12

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Made in lab

<400> 12

gcactccact cttcctttat ttcttaccac

30

<210> 13

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Made in lab

<400> 13

agttacttaa caatgcgcaa ccaaggc

27

<210> 14

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Made in lab

<400> 14

ttgtgtacaa atcatggg

18

<210> 15

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Made in lab

<400> 15

gtaagcccggt cacccatc

18

<210> 16

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Made in lab

<400> 16

gaacaagatg gattgcacgc

20

<210> 17

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Made in lab

<400> 17

gaagaactcg tcaagaaggc

20

INTERNATIONAL SEARCH REPORT

International application No.
PCT/NZ00/00031

A. CLASSIFICATION OF SUBJECT MATTER		
Int. Cl. ⁷ : C07H 21/04, A01H 1/00, C12N 15/82		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) WPIDS, keywords: promoter, regulatory region, pine, pinus. GenBank, EMBL, PDB Nucleic Acids		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Rutledge <i>et al</i> , "Characterization of an AGMOUS homologue from the conifer black spruce (<i>Picea mariana</i>) that produces floral homeotic conversions when expressed in <i>Aridopsis</i> ," <i>The Plant Journal</i> , 1998, 15(5):625-634, [GeneBank Accessions U46582, U69482] whole of document	1-2, 4-7, 9-14, 16-23, 25-30
Y	whole of document	8, 15, 24
Y	Lui J <i>et al</i> , <i>Pinus radiata</i> MADS box protein mRNA, complete cds, Genebank Accession AF023615 sequence	8
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C <input checked="" type="checkbox"/> See patent family annex		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 19 June 2000		Date of mailing of the international search report 27 JUN 2000
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA E-mail address: pct@ipaustalia.gov.au Facsimile No. (02) 6285 3929		Authorized officer GARETH COOK Telephone No : (02) 6283 2541

INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ00/00031

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Taylor B <i>et al</i> , "RNS2: A senescence-associated RNase of <i>Arabidopsis</i> that diverged from S-RNases before speciaiton," <i>Proceedings of the National Acadamy of Science</i> , 1993, 90(11):5118-5122 whole of document	15, 24
P, A	EP 967 278 (Mitsui Chemicals) 29 December 1999 whole of document	1-30

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/NZ00/00031

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report				Patent Family Member	
EP	967 278	AU	36819/99	JP	2000078987
					END OF ANNEX